



## Colonic transit time relates to bacterial metabolism and mucosal turnover in the human gut

Roager, Henrik Munch; Hansen, Lea Benedicte Skov; Bahl, Martin Iain; Frandsen, Henrik Lauritz; Carvalho, Vera; Gøbel, Rikke J.; Dalgaard, Marlene Danner; Sparholt, Morten; Vestergaard, Henrik; Hansen, Torben

*Total number of authors:*  
17

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

### *Citation (APA):*

Roager, H. M., Hansen, L. B. S., Bahl, M. I., Frandsen, H. L., Carvalho, V., Gøbel, R. J., Dalgaard, M. D., Sparholt, M., Vestergaard, H., Hansen, T., Sicheritz-Ponten, T., Nielsen, H. B., Pedersen, O., Lauritzen, L., Kristensen, M., Gupta, R., & Licht, T. R. (2016). *Colonic transit time relates to bacterial metabolism and mucosal turnover in the human gut*. Abstract from 10th INRA-Rowett Symposium: Gut Microbiology, Clermont-Ferrand, France.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Abstract for INRA - ROWETT 2016, 10<sup>th</sup> Joint Symposia - "GUT MICROBIOLOGY: twenty years and counting ..."**

June 20-23, 2016

Clermont-Ferrand, France

### **Colonic transit time relates to bacterial metabolism and mucosal turnover in the human gut**

Henrik M. Roager<sup>1</sup>, Lea B. S. Hansen<sup>2</sup>, Martin I. Bahl<sup>1</sup>, Henrik L. Frandsen<sup>1</sup>, Vera Carvalho<sup>1</sup>, Rikke J. Gøbel<sup>3</sup>, Marlene D. Dalgaard<sup>2</sup>, Morten H. Sparholt<sup>4</sup>, Henrik Vestergaard<sup>3</sup>, Torben Hansen<sup>3</sup>, Thomas S. Pontén<sup>2</sup>, H. Bjørn Nielsen<sup>2</sup>, Oluf Pedersen<sup>3</sup>, Lotte Lauritzen<sup>5</sup>, Mette Kristensen<sup>5</sup>, Ramneek Gupta<sup>2</sup>, Tine R. Licht<sup>1</sup>.

<sup>1</sup>National Food Institute, Technical University of Denmark, Denmark

<sup>2</sup>Department of Systems Biology, Technical University of Denmark, Denmark

<sup>3</sup>The Novo Nordisk Foundation Center for Basic Metabolic Research, Section of Metabolic Genetics, University of Copenhagen, Denmark

<sup>4</sup>Department of Radiology Bispebjerg and Frederiksberg Hospitals

<sup>5</sup>Department of Nutrition, Exercise and Sport, University of Copenhagen, Denmark

Little is known about how colonic transit time relates to human colonic metabolism, and its importance for host health, although stool consistency, a proxy for colonic transit time, has recently been negatively associated with gut microbial richness. To address the relationships between colonic transit time and the gut microbial composition and metabolism, we assessed the colonic transit time of 98 subjects using radiopaque markers, and profiled their gut microbiota by 16S rRNA gene sequencing and their urine metabolome by ultra performance liquid chromatography mass spectrometry. Based on correlation analyses, we show that colonic transit time is associated with overall gut microbial composition, diversity and metabolism. A relatively prolonged colonic transit time associates with high microbial species richness and a shift in colonic metabolism from carbohydrate fermentation to protein catabolism as reflected by microbial metabolites in urine. This results in a number of potentially deleterious protein-derived metabolites. Additionally, longer colonic transit time correlates with metabolites likely reflecting reduced renewal of the colonic mucosa. Together, this suggests that a high gut microbial richness does not *per se* imply a healthy gut microbiota, and contributes to the understanding of the pathophysiology of diseases where increased transit time is a risk factor. Finally, our findings highlight the colonic transit time as an important physiological variable, which should be considered in gut microbiota and metabolomics studies.